

Al Abbar Laboratories for Research and Medical Analysis مختبرات العبارللأبحاث و التحاليل الطبية



إدارة دبن للإعتماد طبقا لمواصفات الأيزو ١٥١٨٩ Dubai Accreditation Department for ISO 15189 المارة دبن للإعتماد طبقا لمواصفات الأيزو

LABORATORY REPORT

Name : ANTHONY KUNYURIA File. No. : AAL02-366117

DOB/Gender

: 27-07-1984 (39 Yrs 4 Month 23 Days/Male)

: Self

Lab No.

: 22233540146

Referral Clinic

Referral Doctor

: Peshawar(Irham Medical Center)

Insurance

Request Date : 20-12-2023 12:58

: No

Clinic File No

IMMUNOLOGY/SEROLOGY/INFECTIOUS DISEASES

Test Name	Result	Units	Ref. Range	Method
**Hepatitis B surface Ag	0.37	Cutoff Index (COI)	Non Reactive: <0.90 Borderline: >= 0.90 to < 1.0 Reactive: >= 1.0	ECLIA
**Hepatitis B "s" Ag Interpretation	Non-Reactive		Non-Reactive	

Full HepB profile is recommended in case of a Reactive result. HBs Ag Index value is an instrument dependant value and should not be used for monitoring.

For monitoring HepB levels (track treatment), a quantitative viral load by PCR is advised.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 20/10/23

Hepatitis C Virus Abs **ECLIA Cutoff Index (COI) Non Reactive: <0.90

Borderline: >= 0.90 to < 1.0

Reactive: >=1.0

Non-Reactive **Hepatitis C Virus Ab Interpretation Non-Reactive

Note: Patient with acute HCV infection may still be non-reactive for HCV abs ('diagnostic window'). Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

ECLIA Cutoff Index (COI) Non Reactive: <1.0 **HIV I & II Abs + p24 Ag 0.17

Reactive: >=1.0

**HIV Interpretation Non-Reactive Non-Reactive

Kindly note above test is just a screening test.

Note: Patients at a very early stage of infection may still be non-reactive for HIV Abs or detect p24 antigen. If HIV is suspected, a control sample is recommended in 4-6 weeks.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

Sample Type: Serum

These tests are accredited under ISO 15189:2012 unless specified by (*) Sample processed on the same day of receipt unless specified otherwise. Test results pertains only the sample tested and to be correlated with clinical history.

Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director

DHA/LS/248469

H/L Collected On: 19-12-2023 18:00:00 Authenticated On: 20-12-2023 14:50:01 Received On: 20-12-2023 12:59:00 Released On: 20-12-2023 15:05:02

Printed On: 20-12-2023 15:31:16

Padmapriya Kumareshan

Sr. Technician

DHA/LS2992011/243822