

LABORATORY REPORT

Name : **ASHRAF ELSAYED ELSAYED ABDELAAL GAABOUB** File. No. : **AAL02-366401**
DOB/Gender : **02-06-1994 (29 Yrs 7 Month 31 Days/Male)** Referral Doctor : **Dr. Enomen Goodluck Ekata**
Lab No. : **22233590064** Referral Clinic : **Peshawar(Irham Medical Center)**
Request Date : **25-12-2023 13:07:13** Clinic File No : **41962**
Insurance : **No**

IMMUNOLOGY/SEROLOGY/INFECTIOUS DISEASES

Test Name	Result	Units	Ref. Range	Method
**Hepatitis B surface Ag	0.46	Cutoff Index (COI)	Non Reactive: <0.90 Borderline: >= 0.90 to < 1.0 Reactive: >= 1.0	ECLIA
**Hepatitis B "s" Ag Interpretation	Non-Reactive		Non-Reactive	

Full HepB profile is recommended in case of a Reactive result. HBs Ag Index value is an instrument dependant value and should not be used for monitoring.

For monitoring HepB levels (track treatment), a quantitative viral load by PCR is advised.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 20/10/23

**Hepatitis C Virus Abs	0.04	Cutoff Index (COI)	Non Reactive: <0.90 Borderline: >= 0.90 to < 1.0 Reactive: >=1.0	ECLIA
**Hepatitis C Virus Ab Interpretation	Non-Reactive		Non-Reactive	

Note: Patient with acute HCV infection may still be non-reactive for HCV abs ('diagnostic window'). Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

**HIV I & II Abs + p24 Ag	0.19	Cutoff Index (COI)	Non Reactive: <1.0 Reactive: >=1.0	ECLIA
**HIV Interpretation	Non-Reactive		Non-Reactive	

Kindly note above test is just a screening test.

Note: Patients at a very early stage of infection may still be non-reactive for HIV Abs or detect p24 antigen. If HIV is suspected, a control sample is recommended in 4-6 weeks.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

Sample Type : Serum

----- End Of Report -----

These tests are accredited under ISO 15189:2012 unless specified by (*)

Sample processed on the same day of receipt unless specified otherwise.

Test results pertains only the sample tested and to be correlated with clinical history.

Reference range related to Age/Gender.

Dr. Solmaz Siddiqui
Laboratory Director
DHA/LS/248469

H/L Collected On : 25-12-2023 10:00:00 Received On : 25-12-2023 13:08:00
Authenticated On : 25-12-2023 14:13:00 Released On : 25-12-2023 15:23:11



Wasik Hasan Tisekar
Laboratory Technician
DHA-P-0039673