

Name : Zuhura Nasoro Hamisi
Lab. No. : 3022119015
Contract. : Peshawer medical L.L.C
Patient No. : 30-83437
File No. :

Sample Date : 08/11/2022 10:42 AM
Report Date : 08/11/2022 14:03 PM
Doctor references : Dr. Tahniyat Iqbal

Branch : Al Borg Lab. Dubai Deirah **Age** : 38 Year **DOB** 08/03/1984 **Sex** : Female

Cholesterol/HDL/LDL/Triglycerides

Test	Result	Unit	Ref. Range
Cholesterol	165	mg/dl	No risk <200 Moderate risk 200 - 240 High risk >240

Comments

Primary sample: Serum
Methodology : Enzymatic colorimetric assay in the presence of peroxidase.

Triglycerides (TG) in Serum	32	mg/dL	Normal<200
-----------------------------	----	-------	------------

Comments

Primary sample: Serum
Methodology : Enzymatic colorimetric assay

HDL Cholesterol	65	mg/dL	Optimum level ≥ 60 Borderline risk: 50-59 High risk: <50 mg/dl
-----------------	----	-------	--

Comments

Primary sample: Serum
Methodology : Homogeneous Enzymatic colorimetric assay.

LDL Cholesterol	94	mg/dl	Optimal <100 Near optimal 100-129 Borderline high 130-159 High 160-189 Very high >189
-----------------	----	-------	---

Comments

Sample type: Serum
Methodology: Calculation

*This is non-accredited test.

Name : Zuhura Nasoro Hamisi
Lab. No. : 3022119015
Contract. : Peshawer medical L.L.C
Patient No. : 30-83437
File No. :

Sample Date : 08/11/2022 10:42 AM
Report Date : 08/11/2022 14:03 PM
Doctor references : Dr. Tahniyat Iqbal

Branch : Al Borg Lab. Dubai Deirah Age : 38 Year DOB 08/03/1984 Sex : Female

Cholesterol/HDL/LDL/Triglycerides

Test	Result	Unit	Ref. Range
Non-HDL Cholesterol	100	mg/dL	very high >220 High 190-219 Borderline high 160-189 Near Ideal 130-159 Ideal for people at risk of heart disease <130 Ideal for people at very high risk of heart disease <100

Comments

Patients with elevated levels of non-HDL-C and normal levels of LDL-C, often have an increased number of LDL particles, increased apo B or increased small, dense LDL particles and these are associated with an increased risk of CVD. Therefore non-HDL-C might be more valuable indicator of cardiovascular risk than LDL-C. The treatment goal for non-HDL cholesterol in persons with high triglycerides (>199 mg/dl) is 30 mg/dl higher than their LDL cholesterol goal.

Reviewed By:



Dr. Shailendra Rathod.MD
Laboratory Director
Physician Specialist Clinical
Pathology
License no. 10944644-001

** End Of Report **