

Name : Naqib Ullah Sample Date : 11/01/2023 15:41 PM
Lab. No. : 3023124531 Report Date : 12/01/2023 13:35 PM
Contract. : Peshawer medical L.L.C Doctor references : Dr. Tahniyat Iqbal
Patient No. : 30-87319
File No. : 30647

Branch : Al Borg Lab. Dubai Age : 28 Year DOB 01/01/1995 Sex : Male
Deirah

Chemistry Unit

Test	Result	Unit	Ref. Range	Lab
T4	5.85	ug/dl	4.87 - 11.72	BUH

Comments

Sample Type: Serum
Methodology: Chemiluminescence immunoassay

Thyroid Stimulating Hormone (TSH)	0.75	uIU/mL	0.27 - 4.2
-----------------------------------	------	--------	------------

Comments

Note: TSH Reference range in pregnant women is varied according to the gestational age:
• First trimester: 0.100-2.500 uIU/mL
• Second trimester: 0.200-3.000 uIU/mL
• Third trimester: 0.300-3.000 uIU/mL
Primary sample: Serum
Methodology: Electrochemiluminescence

Reviewed By:



Dr. Shailendra Rathod.MD
Laboratory Director
Physician Specialist Clinical
Pathology
License no. 10944644-001

BUH : This test was outsourced to Al Borg Medical Laboratories - Abu Dhabi, UAE

**** End Of Report ****

Verified By : Ahmed M. Awad.MD.PhD ,Shailendra Rathod.MD

Printed Date PM 16:15 13/01/2023

Printed By: System

Page 1 Of 1

All Rights Reserved © National Technology

Powered by LDM www.nt-me.com

Name : Naqib Ullah Sample Date : 11/01/2023 15:41 PM
Lab. No. : 3023124531 Report Date : 11/01/2023 17:04 PM
Contract. : Peshawer medical L.L.C Doctor references : Dr. Tahniyat Iqbal
Patient No. : 30-87319
File No. : 30647

Branch : Al Borg Lab. Dubai Age : 28 Year DOB 01/01/1995 Sex : Male
Deirah

Microbiology Unit

Test	Result	Ref. Range
Helicobacter Pylori Stool Antigen (HpSA)	Negative	Negative

Comments

Sample Type: Stool

Methodology: Lateral flow immunoassay

Some drugs can alter intragastric distribution of H. pylori and suppress its activity resulting in false negative results, therefore patient should withhold intake of proton pump inhibitors (PPIs) for a minimum of 2 weeks and antibiotics, metronidazol and bismuth for 4 weeks and H2-receptor antagonists 24 hours prior to testing.

Testing to prove eradication should be performed using a urea breath test or fecal antigen test at least 4 weeks after the completion of therapy.

Malfertheiner P, et al. Current concepts in the management of Helicobacter pylori I infection: the Maastricht III Consensus Report. Gut 2007;56(6):772-81.

Chey WD, et al. ACG Clinical Guideline: Treatment of Helicobacter pylori Infection. Am J Gastroenterol 2017; 112(2):212-239.

Reviewed By:



Dr. Shailendra Rathod.MD
Laboratory Director
Physician Specialist Clinical
Pathology
License no. 10944644-001

**** End Of Report ****

Verified By : Rizvath Noushad

Printed By: System

All Rights Reserved © National Technology

Printed Date PM 16:15 13/01/2023

Page 1 Of 1

Powered by LDM www.nt-me.com