

Name : Muhammad Khalid Arshad
Lab. No. : 3023129209
Contract. : Nas (WN, VN)-Dubai Branch
Patient No. : 30-90546
File No. : 37620

Sample Date : 13/03/2023 08:25 AM
Report Date : 13/03/2023 20:36 PM
Doctor references : Dr. Tahniyat Iqbal

Branch : Al Borg Lab. Dubai Deirah Age : 32 Year DOB 01/01/1991 Sex : Male

Microbiology Unit

Test	Result	Ref. Range
Helicobacter Pylori Stool Antigen (HpSA)	Positive	Negative

Comments

Sample Type: Stool

Methodology: Lateral flow immunoassay

Some drugs can alter intragastric distribution of H. pylori and suppress its activity resulting in false negative results, therefore patient should withhold intake of proton pump inhibitors (PPIs) for a minimum of 2 weeks and antibiotics, metronidazol and bismuth for 4 weeks and H2-receptor antagonists 24 hours prior to testing.

Testing to prove eradication should be performed using a urea breath test or fecal antigen test at least 4 weeks after the completion of therapy.

Malfertheiner P, et al. Current concepts in the management of Helicobacter pylori I infection: the Maastricht III Consensus Report. Gut 2007;56(6):772-81.

Chey WD, et al. ACG Clinical Guideline: Treatment of Helicobacter pylori Infection. Am J Gastroenterol 2017; 112(2):212-239.

Reviewed By:



Dr. Shailendra Rathod.MD
Laboratory Director
Physician Specialist Clinical
Pathology
License no. 10944644-001

**** End Of Report ****

Verified By : Tasneem M Mukri

Printed By: System

All Rights Reserved © National Technology

Printed Date PM 21:46 13/03/2023

Page 1 Of 1

Powered by LDM www.nt-me.com