

Al Abbar Laboratories for Research and مختبرات العبارللأبحاث و التحاليل الطبية Medical Analysis



إدارة دبن للإعتماد طبقا لمواصفات الأيزو ١٥١٨٩ ما Dubai Accreditation Department for ISO 15189

LABORATORY REPORT

Name : PHONE MYAT THU

: 08-05-1997 (26 Yrs 6 Month 2 Days/Male)

Lab No. : **22233270298**

Request Date : 23-11-2023 21:03

Insurance : No

DOB/Gender

File. No. : **AAL02-356308**

Referral Doctor : Dr. Sajid Sanaullah Khan

Referral Clinic : Peshawar(Irham Medical Center)

Clinic File No : 41562

IMMUNOLOGY/SEROLOGY/INFECTIOUS DISEASES

Test Name	Result	Units	Ref. Range	Method
**Hepatitis B surface Ag	0.37	Cutoff Index (COI)	Non Reactive: <0.90 Borderline: >= 0.90 to < 1.0 Reactive: >= 1.0	ECLIA
**Hepatitis B "s" Ag Interpretation	Non-Reactive		Non-Reactive	

Full HepB profile is recommended in case of a Reactive result. HBs Ag Index value is an instrument dependant value and should not be used for monitoring.

For monitoring HepB levels (track treatment), a quantitative viral load by PCR is advised.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 20/10/23

*HCV Abs (serum) 0.05 S/CO Non-Reactive: < 1 CMIA

Reactive: >= 1

**Hepatitis C Virus Ab Interpretation Non-Reactive Non-Reactive

Note: Patient with acute HCV infection may still be non-reactive for HCV abs ('diagnostic window'). Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

**HIV I & II Abs + p24 Ag 0.20 Cutoff Index (COI) Non Reactive: <1.0 ECLIA

Reactive: >=1.0

**HIV Interpretation Non-Reactive Non-Reactive

Kindly note above test is just a screening test.

Note: Patients at a very early stage of infection may still be non-reactive for HIV Abs or detect p24 antigen. If HIV is suspected, a control sample is recommended in 4-6 weeks.

Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 15/10/23

Sample Type : Serum

----- End Of Report -----

These tests are accredited under ISO 15189:2012 unless specified by (*)
Sample processed on the same day of receipt unless specified otherwise.
Test results pertains only the sample tested and to be correlated with clinical history.

Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director DHA/LS/248469 H/L Collected On: 23-11-2023 19:00:00 Received On: 23-11-2023 21:04:00 Authenticated On: 23-11-2023 23:13:15 Released On: 23-11-2023 23:19:18 Printed On: 23-11-2023 23:21:40 Reprinted On: 24-11-2023 09:34:15

agui.

Esmeralda Obenita

DHA/LS/2992011/ 241430

Lab Incharge