biosytech
MEDICAL LABORATORY

BML456492

Laboratory Investigation Report

Name	:	Mr. NARESH GENDA LAL	**Ref No.**	:	38797
DOB	:	03/08/1993	**Sample No.**	:	2408466554
Age / Gender	:	31 Y / Male	**Collected**	:	23/08/2024 13:20
Referred by	:	DR HUMAIRA	**Registered**	:	23/08/2024 21:19
Centre	:	CITICARE MEDICAL CENTER	**Reported**	:	23/08/2024 22:34

BIOCHEMISTRY

Test	Result	Flag	Unit	Reference Range	Methodology
C-REACTIVE PROTEIN (CRP)	**23.4**	**CH**	mg/L	< 5.0 Please note change. Source: Roche IFU.	Particle-enhanced immunoturbidimetric assay

INTERPRETATION NOTES :

1. CRP measurements are used as aid in diagnosis, monitoring, prognosis, and management of suspected inflammatory disorders and associated diseases, acute infections and tissue injury.
2. C-reactive protein is the classic acute phase protein in inflammatory reactions.
3. CRP is the most sensitive of the acute phase reactants and its concentration increases rapidly during inflammatory processes. The CRP response frequently precedes clinical symptoms, including fever. After onset of an acute phase response, the serum CRP concentration rises rapidly and extensively. The increase begins within 6 to 12 hours and the peak value is reached within 24 to 48 hours. Levels above 100 mg/L are associated with severe stimuli such as major trauma and severe infection (sepsis).
4. CRP response may be less pronounced in patients suffering from liver disease.
5. CRP assays are used to detect systemic inflammatory processes (apart from certain types of inflammation such as systemic lupus erythematosus (SLE) and Colitis ulcerosa); to assess treatment of bacterial infections with antibiotics; to detect intrauterine infections with concomitant premature amniorrhexis; to differentiate between active and inactive forms of disease with concurrent infection, e.g. in patients suffering from SLE or Colitis ulcerosa; to therapeutically monitor rheumatic disease and assess anti-inflammatory therapy; to determine the presence of post-operative complications at an early stage, such as infected wounds, thrombosis and pneumonia, and to distinguish between infection and bone marrow transplant rejection.

Sample Type : *Serum*

End of Report

Dr. Adley Mark Fernandes
M.D (Pathology)
Pathologist

Dr. Vyoma V Shah
M.D (Pathology)
Clinical Pathologist

HARSHAD MANIKANDAN
Laboratory Technician
Printed on: 23/08/2024 22:36

This is an electronically authenticated report

P.O Box: 49527 Dubai, UAE Tel: +971 4 398 8567 reports@biosytech.ae www.biosytech.com

biosytech
MEDICAL LABORATORY

BML456492

Laboratory Investigation Report

Name	:	Mr. NARESH GENDA LAL	**Ref No.**	:	38797
DOB	:	03/08/1993	**Sample No.**	:	2408466554
Age / Gender	:	31 Y / Male	**Collected**	:	23/08/2024 13:20
Referred by	:	DR HUMAIRA	**Registered**	:	23/08/2024 21:19
Centre	:	CITICARE MEDICAL CENTER	**Reported**	:	23/08/2024 22:07

HEMATOLOGY

Test	Result	Flag	Unit	Reference Range	Methodology
COMPLETE BLOOD COUNT (CBC)					
HEMOGLOBIN	**12.9**	**L**	g/dL	13.5 - 17.5	Photometric
RBC COUNT	4.5		10^6/µL	4.3 - 5.7	Electrical Impedance
HEMATOCRIT	38.6		%	38 - 50	Calculation
MCV	84.9		fL	82 - 98	Calculation
MCH	28.3		pg	27 - 32	Calculation
MCHC	33.3		g/dL	32 - 37	Calculation
RDW	13.1		%	11.8 - 15.6	Calculation
RDW-SD	38.9		fL		Calculation
MPV	10		fL	7.6 - 10.8	Calculation
PLATELET COUNT	196		10^3/uL	150 - 450	Electrical Impedance
PCT	0.2		%	0.01 - 9.99	Calculation
PDW	17.3		Not Applicable	0.1 - 99.9	Calculation
NUCLEATED RBC (NRBC)^	0.1		/100 WBC		VCS 360 Technology
ABSOLUTE NRBC COUNT^	0.01		10^3/uL		Calculation
EARLY GRANULOCYTE COUNT (EGC)^	0.2		%		VCS 360 Technology
ABSOLUTE EGC^	0		10^3/uL		Calculation
WBC COUNT	9.9		10^3/µL	4 - 11	Electrical Impedance
DIFFERENTIAL COUNT (DC)					
NEUTROPHILS	59		%	40 - 75	VCS 360 Technology
LYMPHOCYTES	32		%	20 - 45	VCS 360 Technology
EOSINOPHILS	3		%	0 - 6	VCS 360 Technology
MONOCYTES	6		%	1 - 6	VCS 360 Technology
BASOPHILS	0		%	0 - 1	VCS 360 Technology
ABSOLUTE COUNT					
ABSOLUTE NEUTROPHIL COUNT	5.7		10^3/uL	1.6 - 8.25	Calculation
ABSOLUTE LYMPHOCYTE COUNT	3.1		10^3/uL	0.8 - 4.95	Calculation
ABSOLUTE MONOCYTE COUNT	0.6		10^3/uL	0.04 - 0.66	Calculation
ABSOLUTE EOSINOPHIL COUNT	0.3		10^3/uL	0 - 0.66	Calculation
ABSOLUTE BASOPHIL COUNT	0.0		10^3/uL	0 - 0.11	Calculation

Dr. Adley Mark Fernandes
M.D (Pathology)
Pathologist

Dr. Vyoma V Shah
M.D (Pathology)
Clinical Pathologist

Reena Babu
Laboratory Technologist

This is an electronically authenticated report

Printed on: 23/08/2024 22:36

P.O Box: 49527 Dubai, UAE Tel: +971 4 398 8567 reports@biosytech.ae www.biosytech.com

biosytech
MEDICAL LABORATORY

BML456492

Laboratory Investigation Report

Name	:	Mr. NARESH GENDA LAL	**Ref No.**	:	38797
DOB	:	03/08/1993	**Sample No.**	:	2408466554
Age / Gender	:	31 Y / Male	**Collected**	:	23/08/2024 13:20
Referred by	:	DR HUMAIRA	**Registered**	:	23/08/2024 21:19
Centre	:	CITICARE MEDICAL CENTER	**Reported**	:	23/08/2024 22:07

HEMATOLOGY

Test	Result	Flag	Unit	Reference Range	Methodology
COMPLETE BLOOD COUNT (CBC)					

INTERPRETATION NOTES : Please note update on CBC report format, reference ranges and method(Beckman Coulter).

Dr. Adley Mark Fernandes
M.D (Pathology)
Pathologist

Dr. Vyoma V Shah
M.D (Pathology)
Clinical Pathologist

Reena Babu
Laboratory Technologist

This is an electronically authenticated report

Printed on: 23/08/2024 22:36

P.O Box: 49527 Dubai, UAE Tel: +971 4 398 8567 reports@biosytech.ae www.biosytech.com

biosytech
MEDICAL LABORATORY

BML456492

Laboratory Investigation Report

Name	:	Mr. NARESH GENDA LAL	**Ref No.**	:	38797
DOB	:	03/08/1993	**Sample No.**	:	2408466554
Age / Gender	:	31 Y / Male	**Collected**	:	23/08/2024 13:20
Referred by	:	DR HUMAIRA	**Registered**	:	23/08/2024 21:19
Centre	:	CITICARE MEDICAL CENTER	**Reported**	:	23/08/2024 22:07

HAEMATOLOGY

Test	Result	Flag	Unit	Reference Range	Methodology
ERYTHROCYTE SEDIMENTATION RATE (ESR)	33	H	mm/hr	< 15 Please note change in reference range and method.	Automated

Comments : Please correlate clinically.

INTERPRETATION NOTES :

Increased ESR is seen in inflammation, pregnancy, anemia, autoimmune disorders (such as rheumatoid arthritis and lupus), infections, some kidney diseases and some cancers (such as lymphoma and multiple myeloma).

The ESR is decreased in polycythemia, hyperviscosity, sickle cell anemia, leukemia, low plasma protein (due to liver or kidney disease), congestive heart failure, hypofibrinogenemia and leukocytosis.

Sample Type : *EDTA Whole Blood*

End of Report

Dr. Adley Mark Fernandes
M.D (Pathology)
Pathologist

Dr. Vyoma V Shah
M.D (Pathology)
Clinical Pathologist

Reena Babu
Laboratory Technologist

This is an electronically authenticated report

Printed on: 23/08/2024 22:36

P.O Box: 49527 Dubai, UAE Tel: +971 4 398 8567 reports@biosytech.ae www.biosytech.com