

# Al Abbar Laboratories for Research and Medical Analysis مختبرات العبار للأبحاث و التجاليل الطبية



إدارة دبن للاعتماد طبقا لمواصفات الأيزو ١٥١٨٩ 15189 Dubai Accreditation Department for ISO 15189

## LABORATORY REPORT

Name : SOUKAINA BENRAQQOUCH

DOB/Gender : 23-12-1992 (31 Yrs 10 Month 20

Days/Female)

Lab No. : 22242990396

Insurance

Request Date : 25-10-2024 22:16:18

Referral Clinic Clinic File No

Referral Doctor

File. No.

: Dr. Humaira Mumtaz

: AAL02-395481

: Citicare Medical Center LLC

: 30914

: NATIONAL GENERAL INSURANCE (HEALTHNET)

#### HORMONES/ENDOCRINE/TUMOR MARKERS

Test Name	Result	Units	Ref. Range	Method
FSH (Follicle Stimulating Hormone)	5.74	mIU/mL	Normal menstruating females: Follicular phase: 3.5-12.5 Ovulation Phase: 4.7-21.5 Luteal phase: 1.7-7.7 Post menopausal: 25.8-134.8	ECLIA

Conversion Formula: (Concentration in mIU/mL) x (1.0) = IU/L.

Determination of the FSH concentration is used in the elucidation of dysfunctions within the hypothalamus pituitary gonads system. The determination of FSH in conjunction with LH is utilized for the following indications: congenital diseases with chromosome aberrations, polycystic ovaries (PCO), amenorrhea (causes),

Depressed gonadotropin levels in men occur in azoospermia. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 09/01/24.

**Luteinising Hormone (LH)** mIU/mL Normal menstruating females: **ECLIA** 17 10

> Follicular phase: 2.4-12.6 Ovulation Phase: 14.0-95.6 Luteal phase: 1.0-11.4 Post menopausal: 7.7-58.5

Conversion Formula: (Concentration in mIU/mL) x (1) = IU/L. Determination of the LH concentration is used in the elucidation of dysfunctions within the hypothalamuspituitarygonads system. The determination of LH in conjunction with FSH is utilized for the following indications: congenital diseases with chromosome aberrations (e.g. Turner's syndrome), polycystic ovaries (PCO), clarifying the causes of amenorrhea, menopausal syndrome, and suspected Leydig cell insufficiency. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration Please note Reference Range Reviewed w.ef 06/02/24.

**TSH (Thyroid Stimulating Hormone)** 0.27-4.20 **ECLIA** 1.5800 uIU/mL

Pregnancy:

1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15

Conversion Formula: (Concentration in IU/ mL) x (1.0) = mIU/L. The determination of TSH serves as the initial test in thyroid diagnostics. Even very slight changes in the concentrations of the free thyroid hormones bring about much greater opposite changes in the TSH level. Accordingly, TSH is a very sensitive

> These tests are accredited under ISO 15189:2012 unless specified by (\*) Sample processed on the same day of receipt unless specified otherwise. Test results pertains only the sample tested and to be correlated with clinical history.

Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director DHA/LS/248469

Received On: 25-10-2024 22:17:00 Patient Sample Collected On: 25-10-2024 18:52:00 Authenticated On: 25-10-2024 23:50:37

Released On: 25-10-2024 23:51:39

Lab Incharge DHA-P-2992011

Esmeralda Obenita







# Al Abbar Laboratories for Research and Medical Analysis مختبرات العبارللأبحاث و التحاليل الطبية



إدارة دبى للإعتماد طبقا لمواصفات الأيزو ١٥١٨٩ - Dubai Accreditation Department for ISO 15189

## LABORATORY REPORT

Name : SOUKAINA BENRAQQOUCH DOB/Gender

: 23-12-1992 (31 Yrs 10 Month 20

Days/Female)

Lab No. : 22242990396

Insurance

Request Date : 25-10-2024 22:16:18

File. No. Referral Doctor : AAL02-395481

: Dr. Humaira Mumtaz

: Citicare Medical Center LLC Referral Clinic

Clinic File No : 30914

: NATIONAL GENERAL INSURANCE (HEALTHNET)

**Test Name** Result Units Ref. Range Method

and specific parameter for assessing thyroid function and is particularly suitable for early detection or exclusion of disorders in the central regulating circuit between the hypothalamus, pituitary and thyroid. Please note Reference Range Reviewed w.ef 15/10/23. Sample Type: Serum

---- End Of Report ----

These tests are accredited under ISO 15189:2012 unless specified by (\*) Sample processed on the same day of receipt unless specified otherwise. Test results pertains only the sample tested and to be correlated with clinical history. Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director DHA/LS/248469

Patient Sample Collected On: 25-10-2024 18:52:00

Authenticated On: 25-10-2024 23:50:37 Released On: 25-10-2024 23:51:39

Received On: 25-10-2024 22:17:00

Esmeralda Obenita

Lab Incharge DHA-P-2992011

