

Al Abbar Laboratories for Research and Medical Analysis مختبرات العبارللأبحاث و التحاليل الطبية



ECLIA

إدارة دبن للاعتماد طبقا لمواصفات الأيزو ١٥١٨٩ 15189 Dubai Accreditation Department for ISO 15189

LABORATORY REPORT

Name : FARIA SABIR SABIR HUSSAIN File. No. : AAL02-424251

DOB/Gender

: 10-07-1996 (28 Yrs 4 Month 30 Days/Female) Referral Doctor

: Dr. Humaira Mumtaz

Lab No.

: 22243130255

: Citicare Medical Center LLC Referral Clinic

Request Date : 08-11-2024 20:50:11

Clinic File No : 43587

Insurance : No

HORMONES/ENDOCRINE/TUMOR MARKERS

Test Name Result Units Ref. Range Method

FSH (Follicle Stimulating Hormone) 15.00 mIU/mL Normal menstruating females:

Follicular phase: 3.5-12.5 Ovulation Phase: 4.7-21.5 Luteal phase: 1.7-7.7

Post menopausal: 25.8-134.8

Conversion Formula: (Concentration in mIU/mL) x (1.0) = IU/L.

Determination of the FSH concentration is used in the elucidation of dysfunctions within the hypothalamus pituitary gonads system. The determination of FSH in conjunction with LH is utilized for the following indications: congenital diseases with chromosome aberrations, polycystic ovaries (PCO), amenorrhea (causes), and menopausal syndrome.

Depressed gonadotropin levels in men occur in azoospermia. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration.

Please note Reference Range Reviewed w.ef 09/01/24.

mIU/mL **ECLIA Luteinising Hormone (LH)** 75.10 Normal menstruating females:

Follicular phase: 2.4-12.6 Ovulation Phase: 14.0-95.6 Luteal phase: 1.0-11.4 Post menopausal: 7.7-58.5

Conversion Formula: (Concentration in mIU/mL) x (1) = IU/L. Determination of the LH concentration is used in the elucidation of dysfunctions within the hypothalamuspituitarygonads system. The determination of LH in conjunction with FSH is utilized for the following indications: congenital diseases with chromosome aberrations (e.g. Turner's syndrome), polycystic ovaries (PCO), clarifying the causes of amenorrhea, menopausal syndrome, and suspected Leydig cell insufficiency. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. > 5 mg/day) until at least 8 hours following the last biotin administration Please note Reference Range Reviewed w.ef 06/02/24 .

TSH (Thyroid Stimulating Hormone) µIU/mL 0.27-4.20 **ECLIA**

Pregnancy:

1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15

Conversion Formula: (Concentration in IU/ mL) x (1.0) = mIU/L. The determination of TSH serves as the initial test in thyroid diagnostics. Even very slight changes in the concentrations of the free thyroid hormones bring about much greater opposite changes in the TSH level. Accordingly, TSH is a very sensitive

> These tests are accredited under ISO 15189:2012 unless specified by (*) Sample processed on the same day of receipt unless specified otherwise. Test results pertains only the sample tested and to be correlated with clinical history. Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director

DHA/LS/248469

Patient Sample Collected On: 07-11-2024 10:29:00 Authenticated On: 08-11-2024 21:55:34

Received On: 08-11-2024 20:50:00 Released On: 08-11-2024 21:58:07

Lab Incharge DHA-P-2992011

Esmeralda Obenita





Al Abbar Laboratories for Research and مختبرات العبارللأبحاث و التحاليل الطبية Medical Analysis



إدارة دبن للإعتماد طبقا لمواصفات الأيزو ١٥١٨٩ ما المام Dubai Accreditation Department for ISO 15189

LABORATORY REPORT

Name : FARIA SABIR SABIR HUSSAIN

: 10-07-1996 (28 Yrs 4 Month 30 Days/Female) Referral Doctor

Lab No. : **22243130255**

Request Date : 08-11-2024 20:50:11

Insurance : No

DOB/Gender

File. No. : **AAL02-424251**

Referral Doctor : Dr. Humaira Mumtaz

Referral Clinic : Citicare Medical Center LLC

Clinic File No : 43587

Test Name Result Units Ref. Range Method

and specific parameter for assessing thyroid function and is particularly suitable for early detection or exclusion of disorders in the central regulating circuit

between the hypothalamus, pituitary and thyroid. Please note Reference Range Reviewed w.ef 15/10/23.

Sample Type : Serum

These tests are accredited under ISO 15189:2012 unless specified by (*)
Sample processed on the same day of receipt unless specified otherwise.

Test results pertains only the sample tested and to be correlated with clinical history.

Reference range related to Age/Gender.

Dr. Solmaz Siddiqui Laboratory Director DHA/LS/248469 Patient Sample Collected On: 07-11-2024 10:29:00

Authenticated On: 08-11-2024 21:55:34

Received On: 08-11-2024 20:50:00

Released On: 08-11-2024 21:58:07 Lab Incharge DHA-P-2992011



Esmeralda Obenita Lab Incharge

