





Mr. EMMANUEL NIJI

PID NO: 46832

Age: 45 Years Sex: Male

Sample Collected At :

CITICARE MEDICAL CENTER

Unit G03, Al Barsha South Bldg, Al Barhsa South

Reference: CITICARE MEDICAL CENTER

Third, Dubai

VID: 5050105067

Registered on:

13-May-2025 05:38 PM

Collected on :

13-May-2025 10:47 AM

Reported on:

13-May-2025 07:04 PM

<u>Investigation</u> <u>Observed Value</u> <u>Flag Unit</u> <u>Biological Reference Interval</u>

* HEPATITIS B SURFACE ANTIGEN (HBSAG)

* HIV I & II ANTIBODY AND P24 ANTIGEN

(Serum, ECLIA)

0.336 COI Non-Reactive: < 0.9

Borderline: = or > 0.9 - < 1.0

Reactive: = or > 1.0

Note changes in method and

reference range. Source: Roche IFU.

INTERPRETATION:

• A positive HBsAg test result means that the patient is infected with acute or chronic hepatitis B virus or chronic HBV carrier state.

0.215

• A negative result implies the patient is not infected with hepatitis B.

(Serum, ECLIA)

S/CO Non-Reactive: < 1.0
Reactive: = or > 1.0

Source: Roche IFU.

INTERPRETATION:

- A negative test result does not completely rule out the possibility of an infection with HIV. Serum or plasma samples from the very early (preseroconversion) phase or the late phase of HIV infection can occasionally
- yield negative findings. Yet unknown HIV variants can also lead to a negative HIV finding. The presence of antibodies to HIV is not a diagnosis of AIDS.

 For diagnostic purposes, the results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.
- This is a screening test.

References:

Roche Cobas IFU.

----- End Of Report -----

ayona V. Shah

DR. ADLEY MARK FERNANDES M.D (Pathology) Pathologist DR. VYOMA SHAH M.D (Pathology) Clinical Pathologist

Page 1 of 1

NEZAR ALI

Laboratory Technologist

This is an Electronically Authenticated Report.

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189 unless specified by (*).





