



Ms. AYELA PYAE

PID NO: 45782

Age: 22 Years Sex: Female 11-Aug-2002



Reference: Dr. AMAIZAH ISHTIAQ

Sample Collected At:

CITICARE MEDICAL CENTER

Unit G03, Al Barsha South Bldg, Al Barhsa South

Third, Dubai

VID: 5050111559

Registered on:

31-May-2025 08:41 PM

Collected on:

31-May-2025 12:45 PM

Reported on :

31-May-2025 09:49 PM

<u>Investigation</u>	Observed Value	<u>Flag</u>	<u>Unit</u>	Biological Reference Int	<u>terval</u> <u>Method</u>
COMPLETE BLOOD COUNT (CBC)					
HEMOGLOBIN	11.3	L	g/dL	12 - 15.5	Photometric
RBC COUNT	4.3		10^6/μL	3.9 - 5	Electrical Impedance
HEMATOCRIT	35.0		%	35 - 45	Calculation
MCV	81.0	L	fL	82 - 98	Calculation
мсн	26.1	L	pg	27 - 32	Calculation
мснс	32.2		g/dL	32 - 37	Calculation
* RDW	13.5		%	11.9 - 15.5	Calculation
* RDW-SD	38.10		fL		Calculation
MPV	8.3		fL	7.6 - 10.8	Calculation
PLATELET COUNT	345		10^3/uL	150 - 450	Electrical Impedance
* NUCLEATED RBC (NRBC)	0.20		/100 WBC		VCS 360 Technology
* ABSOLUTE NRBC COUNT	0.02		10^3/uL		Calculation
<b>TOTAL &amp; DIFFERENTIAL COUNT (DC)</b>					
WBC COUNT	11.0		10^3/μL	4 - 11	Electrical Impedance
NEUTROPHILS	49		%	40 - 75	VCS 360 Technology
LYMPHOCYTES	46		%	30 - 60	VCS 360 Technology
EOSINOPHILS	1		%	0 - 6	VCS 360 Technology
MONOCYTES	4		%	1 - 6	VCS 360 Technology
BASOPHILS	0		%	0 - 1	VCS 360 Technology
ABSOLUTE COUNT					
ABSOLUTE NEUTROPHIL COUNT	5.4		10^3/uL	1.6 - 8.25	Calculation
ABSOLUTE LYMPHOCYTE COUNT	5.1		10^3/uL	1.2 - 6.6	Calculation
ABSOLUTE MONOCYTE COUNT	0.4		10^3/uL	0.04 - 0.66	Calculation
ABSOLUTE EOSINOPHIL COUNT	0.1		10^3/uL	0 - 0.66	Calculation
ABSOLUTE BASOPHIL COUNT	0		10^3/uL	0 - 0.11	Calculation

ayana V. Shah

DR. ADLEY MARK FERNANDES M.D (Pathology) Pathologist

Sample Type: EDTA Whole Blood

DR. VYOMA SHAH M.D (Pathology) Clinical Pathologist

Laboratory Technologist

JALADINI DULANKA

31-May-2025 10:51 PM

This is an Electronically Authenticated Report.

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189 unless specified by (\*).













Ms. AYELA PYAE

PID NO: 45782

Age: 22 Years Sex: Female 11-Aug-2002



Reference: Dr. AMAIZAH ISHTIAQ

Sample Collected At:

CITICARE MEDICAL CENTER

Unit G03, Al Barsha South Bldg, Al Barhsa South

Third, Dubai

VID: 5050111559

Registered on:

31-May-2025 08:41 PM

Collected on:

31-May-2025 12:45 PM

Reported on :

31-May-2025 10:49 PM

<u>Investigation</u> <u>Observed Value</u> <u>Flag</u> <u>Unit</u> <u>Biological Reference Interval</u>

\* C-REACTIVE PROTEIN (CRP)

(Serum, Particle-enhanced immunoturbidimetric assay)

**5.19** H mg/L < 5.0

Please note change. Source: Roche IFU.

## INTERPRETATION:

- CRP measurements are used as aid in diagnosis, monitoring, prognosis, and management of suspected inflammatory disorders and associated diseases, acute infections and tissue injury.
- C-reactive protein is the classic acute phase protein in inflammatory reactions.
- CRP is the most sensitive of the acute phase reactants and its concentration increases rapidly during inflammatory processes. The CRP response frequently precedes clinical symptoms, including fever. After onset of an acute phase response, the serum CRP concentration rises rapidly and extensively. The increase begins within 6 to 12 hours and the peak value is reached within 24 to 48 hours. Levels above 100 mg/L are associated with severe stimuli such as major trauma and severe infection (sepsis).
- CRP response may be less pronounced in patients suffering from liver disease.
- CRP assays are used to detect systemic inflammatory processes (apart from certain types of inflammation such as systemic lupus erythematosus (SLE) and Colitis ulcerosa); to assess treatment of bacterial infections with antibiotics; to detect intrauterine infections with concomitant premature amniorrhexis; to differentiate between active and inactive forms of disease with concurrent infection, e.g. in patients suffering from SLE or Colitis ulcerosa; to therapeutically monitor rheumatic disease and assess anti-inflammatory therapy; to determine the presence of post-operative complications at an early stage, such as infected wounds, thrombosis and pneumonia, and to distinguish between infection and bone marrow transplant rejection."

ayana V. Shah

DR. ADLEY MARK FERNANDES M.D (Pathology) Pathologist DR. VYOMA SHAH M.D (Pathology) Clinical Pathologist

Printed on:

Laboratory Technician

HARSHAD MANIKANDAN

31-May-2025 10:51 PM

STATION WITH THE PARTY OF THE P





Page 2 of 2

This is an Electronically Authenticated Report.

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189 unless specified by (\*).