



Ms. YASODA RAJAGOPAL.

PID NO:

Age: 57 Years Sex: Female DOB: 03-May-1968



Reference: Dr. AMAIZAH ISHTIAQ

Sample Collected At:

CITICARE MEDICAL CENTER

Unit G03, Al Barsha South Bldg, Al Barhsa South

Third, Dubai

VID: 5060103137

Registered on:

12-Jun-2025 12:59 PM

Collected on:

11-Jun-2025 05:00 PM

Reported on:

12-Jun-2025 03:29 PM

					_
<u>Investigation</u>	Observed Value	<u>Flag</u>	<u>Unit</u>	Biological Reference Int	erval <u>Method</u>
COMPLETE BLOOD COUNT (CBC)					
HEMOGLOBIN	11.3	L	g/dL	12 - 15.5	Photometric
RBC COUNT	4.7		10^6/μL	3.9 - 5	Electrical Impedance
HEMATOCRIT	35.1		%	35 - 45	Calculation
MCV	75.5	L	fL	82 - 98	Calculation
МСН	24.3	L	pg	27 - 32	Calculation
мснс	32.1		g/dL	32 - 37	Calculation
* RDW	12.7		%	11.9 - 15.5	Calculation
* RDW-SD	34.1		fL		Calculation
MPV	10.7		fL	7.6 - 10.8	Calculation
PLATELET COUNT	175		10^3/uL	150 - 450	Electrical Impedance
* NUCLEATED RBC (NRBC)	0.0		/100 WBC		VCS 360 Technology
* ABSOLUTE NRBC COUNT	0.00		10^3/uL		Calculation
TOTAL & DIFFERENTIAL COUNT (DC)					
WBC COUNT	11.3	Н	10^3/μL	4 - 11	Electrical Impedance
NEUTROPHILS	70		%	40 - 75	VCS 360 Technology
LYMPHOCYTES	24	L	%	30 - 60	VCS 360 Technology
EOSINOPHILS	2		%	0 - 6	VCS 360 Technology
MONOCYTES	4		%	1 - 6	VCS 360 Technology
BASOPHILS	0		%	0 - 1	VCS 360 Technology
ABSOLUTE COUNT					
ABSOLUTE NEUTROPHIL COUNT	7.9		10^3/uL	1.6 - 8.25	Calculation
ABSOLUTE LYMPHOCYTE COUNT	2.7		10^3/uL	1.2 - 6.6	Calculation
ABSOLUTE MONOCYTE COUNT	0.5		10^3/uL	0.04 - 0.66	Calculation
ABSOLUTE EOSINOPHIL COUNT	0.2		10^3/uL	0 - 0.66	Calculation
ABSOLUTE BASOPHIL COUNT	0		10^3/uL	0 - 0.11	Calculation

na V. Shah

DR. ADLEY MARK FERNANDES M.D (Pathology) Pathologist

Sample Type: EDTA Whole Blood

DR. VYOMA SHAH M.D (Pathology) **Clinical Pathologist**

THAHSINA ANEES Laboratory Technologist

This is an Electronically Authenticated Report.

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189 unless specified by (*).







Page 1 of 2







Ms. YASODA RAJAGOPAL.

PID NO:

Age: 57 Years Sex: Female

DOB: 03-May-1968

Reference: Dr. AMAIZAH ISHTIAQ

Sample Collected At:

CITICARE MEDICAL CENTER

Unit G03, Al Barsha South Bldg, Al Barhsa South

Third, Dubai

VID: 5060103137

Registered on:

12-Jun-2025 12:59 PM

Collected on: 11-Jun-2025 05:00 PM

Reported on:

12-Jun-2025 03:24 PM

Investigation Observed Value Flag <u>Unit</u> **Biological Reference Interval**

8.83

* C-REACTIVE PROTEIN (CRP)

(Serum, Particle-enhanced immunoturbidimetric assay)

< 5.0 Н mg/L

> Please note change. Source: Roche IFU.

INTERPRETATION:

- CRP measurements are used as aid in diagnosis, monitoring, prognosis, and management of suspected inflammatory disorders and associated diseases, acute infections and tissue injury.
- C-reactive protein is the classic acute phase protein in inflammatory reactions.
- CRP is the most sensitive of the acute phase reactants and its concentration increases rapidly during inflammatory processes. The CRP response frequently precedes clinical symptoms, including fever. After onset of an acute phase response, the serum CRP concentration rises rapidly and extensively. The increase begins within 6 to 12 hours and the peak value is reached within 24 to 48 hours. Levels above 100 mg/L are associated with severe stimuli such as major trauma and severe infection (sepsis).
- CRP response may be less pronounced in patients suffering from liver disease.
- CRP assays are used to detect systemic inflammatory processes (apart from certain types of inflammation such as systemic lupus erythematosus (SLE) and Colitis ulcerosa); to assess treatment of bacterial infections with antibiotics; to detect intrauterine infections with concomitant premature amniorrhexis; to differentiate between active and inactive forms of disease with concurrent infection, e.g. in patients suffering from SLE or Colitis ulcerosa; to therapeutically monitor rheumatic disease and assess anti-inflammatory therapy; to determine the presence of post-operative complications at an early stage, such as infected wounds, thrombosis and pneumonia, and to distinguish between infection and bone marrow transplant rejection."

DR. ADLEY MARK FERNANDES M.D (Pathology) **Pathologist**

DR. VYOMA SHAH M.D (Pathology) **Clinical Pathologist**

ELOISA MAY DELMO Laboratory Technologist

This is an Electronically Authenticated Report.

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189 unless specified by (*).







Page 2 of 2