

Date:10/08/2025	Patient Name:	JENNETH LOGRONIO	Patient ID:45727
Gender: F	Age: 42Y	Ref Physician: DR. RAHMAN	

## **Abdominal and Pelvis Ultrasonography Examination**

- The liver is normal in size with mild increased parenchymal echogenicity. Intrahepatic and extrahepatic biliary ducts are normal, no evidence of mass lesion.
   Right lobe- 147mm (N= 120-160mm)
   Left lobe- 80mm (N= 100 mm)
- The gallbladder is well-distended and appears normal in size. There is a 69x21mm in size tubular shape non-shadowing polypoid mass lesion almost occupying the entire gallbladder lumen. There is also a 10x7mm in size non-mobile non-shadowing polypoid nodule attached to the gallbladder wall. No evidence of wall thickening, lithiasis or pericholecystic fluid collection.
- Pancreas is normal in size with homogenous parenchymal echotexture, no evidence of calcification, focal lesion or peripancreatic fluid collection.
- Spleen is normal in size with homogenous parenchymal echotexture, no evidence of mass lesion.
- Both kidneys are normal in location, size and parenchymal echogenicity. Cortico-medullary differentiation is preserved, perinephric planes appear normal, no evidence of mass lesion. There are 3mm in sizes non-obstructive stones in both kidneys.
   Right kidney- 120x50mm (cortex- 16mm)
  - Left kidney 123x49mm (cortex-16mm)
- The urinary bladder wall thickness is normal, no evidence of diverticule or abnormal mass lesion in the wall and within the urinary bladder. No calculi seen.
   Urinary bladder volume=558ml
  - **Urinary bladder Post voiding volume=32ml (5%)**
- The uterus is normal in size and volume with homogenous parenchymal echotextute and endometrial thickness of 10mm. No evidence of mass lesion.
- Both ovaries appear normal, no evidence of mass lesion.
- No evidence of free fluid within the abdominal cavity and pelvic region.
- No evidence of bowel wall thickening, intussusceptions or any sign for acute appendicitis in this
  examination.

## **IMPRESSION:**

- The liver is normal in size with mild increased parenchymal echogenicity (mild fatty infiltration).
- There is a 69x21mm in size tubular shape non-shadowing polypoid mass lesion almost occupying the entire gallbladder lumen.
- There is also a 10x7mm in size non-mobile non-shadowing polypoid nodule attached to the gallbladder wall.
- There are 3mm in sizes non-obstructive stones in both kidneys.

Please correlate with clinical and laboratory findings. Thank you very much.

Dr. Eliass Bohlouli Kery Bozorg Specialist Radiology License No.: 00267151-001 مرخـز سيتيكير الطبي ذم م CITICARE MEDICAL CENTER LLC

DR. Eliass Bohlouli Kery Bozorg Specialist Radiologist

04 297 4343

www.citicare24.ae

infociticaie24.ae

04 - 77 00948

•97 1 5ó 933 8664

Al Salam Building, Al Barsha South, Allan Near Mirarle