

Patient Name	: Ms. MARIA YANEY PEREZ GOMEZ	Sample UID No.	: 4106975
Age / Gender	: 47 Y / Female	Sample Collected On	: 09-08-2025 16:45
Patient ID	: QLD106517	Registered On	: 09-08-2025 22:08
Referred By	: Dr. AMAIZAH	Reported on	: 10-08-2025 08:26
Referral Client	: CITICARE MEDICAL CENTER	External Patient ID	: 38080
Emirates ID / Passport No	:	Print Version	: V.1

Department of BIOCHEMISTRY

Investigation	Results	Flag	Units	Biological Reference Interval	Method
* C-REACTIVE PROTEIN (CRP)	6.2	H	mg/L	< 5	Particle enhanced immunoturbidimetric assay

Sample: Serum

Comments :

CLINICAL IMPLICATIONS:

1. CRP is the most sensitive acute phase reactant that can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery or neoplastic proliferation. CRP levels may predict future cardiovascular events and can be used as a screening tool.
2. The traditional test of CRP has added significance over the elevated ESR, which may be influenced by altered physiologic states. CRP tends to increase before rises in antibody titres and ESR level occurs. CRP levels also tend to decrease sooner than ESR levels.
3. The traditional test for CRP is elevated in rheumatic fever, RA, myocardial infarction, malignancy, bacterial and viral infections. The positive test indicates active inflammation but not its cause. In RA, the traditional test for CRP becomes negative with successful treatment and indicates that the inflammation has subsided.
4. High sensitive measurement of CRP (hs-CRP) are useful in assessing vascular inflammation and cardiovascular stratification. A single test for hs-CRP may not reflect an individual patient basal hs-CRP level, therefore follow up tests or serial measurements may be required in patients presenting with increased hs-CRP levels.

INTERFERING FACTORS: Haemolysed or lipemic sample may alter the results.

REFERENCE:

- 1) Manual of Laboratory and Diagnostics -Frances Fischbach Marshall B. Dunning III [9th Edition]
- 2) Tietz clinical guide to Laboratory tests(Fourth edition) ALAN H.B.WU

Note:

"The analytes with asterisk (*) symbol are non-accredited parameters."
"QLabs compliance with ISO 15189:2022 standards"

Maqsood Rahman

Maqsood Rahman
Lab Technologist

DHA No:48036476-001



Dr. Vidhya Mohan

Dr. Vidhya Mohan
Specialist Clinical Pathologist
Clinical Pathologist
DHA No. 23553203-004

Dr. Dheepa Manoharan

Dr. Dheepa Manoharan
Medical Director
Specialist Microbiologist
DHA No. 00231751-004

Patient Name	: Ms. MARIA YANEY PEREZ GOMEZ	Sample UID No.	: EB4106975
Age / Gender	: 47 Y / Female	Sample Collected On	: 09-08-2025 16:45
Patient ID	: QLD106517	Registered On	: 09-08-2025 22:08
Referred By	: Dr. AMAIZAH	Reported on	: 10-08-2025 08:26
Referral Client	: CITICARE MEDICAL CENTER	External Patient ID	: 38080
Emirates ID / Passport No	:	Print Version	: V.1

Department of HEMATOLOGY

COMPREHENSIVE COMPLETE BLOOD COUNT

<u>Investigation</u>	<u>Results</u>	<u>Flag</u>	<u>Units</u>	<u>Biological Reference Interval</u>	<u>Method</u>
HEMOGLOBIN	9.7	L	g/dl	12-15	photometric
RBC COUNT	3.48	L	10 ⁶ /uL	3.8-4.8	Electrical Impedance
HEMATOCRIT	29.8	L	%	37-47	Calculation
MCV	85.5		fL	78-100	Calculation
MCH	27.9		pg	27-31	Calculation
MCHC	32.7		g/dl	31-35	Calculation
RDW	16.4	H	%	9.3-16	Calculation
RDW-SD	47.7		fL	38.9-49	Calculation
MPV	10.5		fL	8.8-12.5	Calculation
PLATELET COUNT	301		10 ³ /uL	150-400	Electrical Impedance
* PCT	0.3		%	0.01-9.99	Calculation
* PDW	17.4			0.1-99.9	Calculation
* NUCLEATED RBC (NRBC)^	0.3		/100 WBC		Flow Cytometry
* ABSOLUTE NRBC COUNT^	0.02		10 ³ /uL		Calculation
* EARLY GRANULOCYTE COUNT (EGC)^	0.11		%		Flow Cytometry
* ABSOLUTE EGC^	0.01		10 ³ /uL		Calculation
WBC COUNT	6.5		10 ³ /uL	4-11	Electrical Impedance
* Neutrophil	63.27		%	40-80	VCS-Method
* Lymphocyte	24.55		%	20-40	VCS-Method
* Eosinophil	1.99		%	1-8	VCS-Method
* Monocyte	9.29		%	2-10	VCS-Method
* Basophil	0.9		%	0-2	VCS-Method
* ABSOLUTE NEUTROPHIL COUNT	4.09		10 ³ /uL	1.5-7	Calculation
* ABSOLUTE LYMPHOCYTE COUNT	1.59		10 ³ /uL	1.5-4	Calculation
* ABSOLUTE MONOCYTE COUNT	0.6		10 ³ /uL	0-0.8	Calculation
* ABSOLUTE EOSINOPHIL COUNT	0.13		10 ³ /uL	0-0.6	Calculation
* ABSOLUTE BASOPHIL COUNT	0.06		10 ³ /uL	0-0.2	Calculation

Sample: EDTA Whole Blood

Note:

"The analytes with asterisk (*) symbol are non-accredited parameters."

"QLabs compliance with ISO 15189:2022 standards"



Mohammed Jahfar Kuttikkattil
Lab Technologist

DHA No:05143389-001





Dr. Vidhya Mohan
Specialist Clinical Pathologist
Clinical Pathologist
DHA No. 23553203-004

Page 2 of 3

Patient Name	: Ms. MARIA YANEY PEREZ GOMEZ	Sample UID No.	: 4106975
Age / Gender	: 47 Y / Female	Sample Collected On	: 09-08-2025 16:45
Patient ID	: QLD106517	Registered On	: 09-08-2025 22:08
Referred By	: Dr. AMAIZAH	Reported on	: 10-08-2025 02:24
Referral Client	: CITICARE MEDICAL CENTER	External Patient ID	: 38080
Emirates ID / Passport No	:	Print Version	: V.1

Department of IMMUNOLOGY

<u>Investigation</u>	<u>Results</u>	<u>Flag</u>	<u>Units</u>	<u>Biological Reference Interval</u>	<u>Method</u>
IGE TOTAL ANTIBODY	72.7		IU/mL	0 - 100	ECLIA

Sample: Serum

Interpretation Notes :

An elevated level of total IgE indicates an allergic process is likely present, but it will not indicate what a person is allergic to. In general, the greater the number of things a person is allergic to, the higher the total IgE level may be. An IgE elevation can also indicate the presence of a parasitic infection but cannot be used to determine the type of infection.

A normal IgE level makes it less likely that a person has allergies but does not rule them out due to the length of time between exposures. In between exposures, a person's IgE level may drop.

- END OF REPORT -

"QLabs compliance with ISO 15189:2022 standards"

Maqsood Rahman

Maqsood Rahman
Lab Technologist

DHA No:48036476-001



Dr. Vidhya Mohan

Dr. Vidhya Mohan
Specialist Clinical Pathologist
Clinical Pathologist
DHA No. 23553203-004

Dr. Dheepa Manoharan

Dr. Dheepa Manoharan
Medical Director
Specialist Microbiologist
DHA No. 00231751-004