



CITICARE

Your Health is our Priority

We Promise We Care

Date:24/08/2025	Patient Name: RIZALINA ACOSTA	Patient ID:45724
Gender: F	Age: 42Y	Ref Physician : DR. KEERTHANA

Abdominal and Pelvis Ultrasonography Examination

- The liver is normal in size and parenchymal echogenicity. Intrahepatic and extrahepatic biliary ducts are normal, no evidence of mass lesion.
Right lobe- 130mm (N= 120-160mm) Left lobe- 71mm (N= 100 mm)
- The gallbladder is well-distended and appears normal in size. There is a 8mm in size echogenic stone having posterior acoustic shadow within the lumen of gallbladder. No evidence of wall thickening, polyp or pericholecystic fluid collection.
- Pancreas is normal in size with homogenous parenchymal echotexture, no evidence of calcification, focal lesion or peripancreatic fluid collection.
- Spleen is normal in size with homogenous parenchymal echotexture, no evidence of mass lesion.
- Both kidneys are normal in location, size and parenchymal echogenicity. Cortico-medullary differentiation is preserved, perinephric planes appear normal, no evidence of mass lesion. There are 3mm in sizes non-obstructive stones in both kidneys. The pelvicalyceal systems and ureters appear normal.
Right kidney- 100x39mm (cortex- 14mm)
Left kidney - 104x41mm (cortex-14mm)
- The urinary bladder wall thickness is normal, no evidence of diverticule or abnormal mass lesion in the wall and within the urinary bladder. No calculi seen.
Urinary bladder pre-voiding volume=363ml
Urinary bladder post voiding volume=36ml (10%)
- The uterus is normal in size and volume with endometrial thickness of 10mm, no evidence of mass lesion. There are 5-6mm in sizes well-defined anechoic cysts in the peripheral region of both ovaries.
- There are minimal amount of anechoic free fluid in the posterior cul-de-sac region.
- No evidence of bowel wall thickening, intussusceptions or any sign for acute appendicitis in this examination.

IMPRESSION:

- There is a 8mm in size echogenic stone having posterior acoustic shadow within the lumen of gallbladder (cholecystolithiasis).
- There are 3mm in sizes non-obstructive stones in both kidneys.
- There are 5-6mm in sizes well-defined anechoic cysts in the peripheral region of both ovaries (polycystic ovaries).
- There are minimal amount of anechoic free fluid in the posterior cul-de-sac region.

Dr. Eliass Bohlouli Kery Bozorg
Specialist Radiology
License No.: 00267151-001
مركز سيتيكير الطبي
CITICARE MEDICAL CENTER LLC

Please correlate with clinical and laboratory findings.
Thank you very much.

DR. Eliass Bohlouli Kery Bozorg
Specialist Radiologist

04 297 4343

www.citicare24.ae



infociticaie24.ae

04-77 00948

•971 56 933 8664

Al Salam Building, Al Barsha South, Allan Near Mirarle