

Name : Hemalatha Jinnagara **Sample Date** : 19/08/2022 20:26 PM
Lab. No. : 3022111678 **Report Date** : 19/08/2022 20:54 PM
Contract. : Peshawer medical L.L.C
Patient No. : 30-78341
File No. : 34343

Branch : Al Borg Lab. Dubai **Age** : 42 Year **DOB** 22/05/1980 **Sex** : Female
Deirah

Cholesterol/HDL/LDL/Triglycerides

Test	Result	Unit	Ref. Range
Cholesterol	208 H	mg/dl	No risk <200 Moderate risk 200 - 240 High risk >240

Comments

Primary sample: Serum
 Methodology : Enzymatic colorimetric assay in the presence of peroxidase.

Triglycerides (TG) in Serum	162	mg/dL	Normal<200
-----------------------------	-----	-------	------------

Comments

Primary sample: Serum
 Methodology : Enzymatic colorimetric assay

HDL Cholesterol	33 L	mg/dL	Optimum level ≥ 60 Borderline risk: 50-59 High risk: <50 mg/dl
-----------------	------	-------	--

Comments

Primary sample: Serum
 Methodology : Homogeneous Enzymatic colorimetric assay.

LDL Cholesterol	142 H	mg/dl	Optimal <100 Near optimal 100-129 Borderline high 130-159 High 160-189 Very high >189
-----------------	-------	-------	---

Comments

Sample type: Serum
 Methodology: Calculation

*This is non-accredited test.

Name : Hemalatha Jinnagara **Sample Date** : 19/08/2022 20:26 PM
Lab. No. : 3022111678 **Report Date** : 19/08/2022 20:54 PM
Contract. : Peshawer medical L.L.C
Patient No. : 30-78341
File No. : 34343

Branch : Al Borg Lab. Dubai Deirah **Age** : 42 Year **DOB** 22/05/1980 **Sex** : Female

Cholesterol/HDL/LDL/Triglycerides

Test	Result	Unit	Ref. Range
Non-HDL Cholesterol	175 H	mg/dL	very high >220 High 190-219 Borderline high 160-189 Near Ideal 130-159 Ideal for people at risk of heart disease <130 Ideal for people at very high risk of heart disease <100

Comments

Patients with elevated levels of non-HDL-C and normal levels of LDL-C, often have an increased number of LDL particles, increased apo B or increased small, dense LDL particles and these are associated with an increased risk of CVD. Therefore non-HDL-C might be more valuable indicator of cardiovascular risk than LDL-C.
 The treatment goal for non-HDL cholesterol in persons with high triglycerides (>199 mg/dl) is 30 mg/dl higher than their LDL cholesterol goal.

Reviewed By:



Dr. Shailendra Rathod.MD
 Laboratory Director
 Physician Specialist Clinical Pathology
 License no. 10944644-001

** End Of Report **

Name : Hemalatha Jinnagara
Lab. No. : 3022111678
Contract. : Peshawer medical L.L.C
Patient No. : 30-78341
File No. : 34343

Sample Date : 19/08/2022 20:26 PM
Report Date : 19/08/2022 20:49 PM

Branch : Al Borg Lab. Dubai Deirah Age : 42 Year DOB 22/05/1980 Sex : Female

Chemistry Unit

Test	Result	Unit	Ref. Range
Glucose in Plasma (Fasting)	96	mg/dl	70 - 100 Normal: 70 - 100 Pre-Diabetes: 100 -125 Diabetes: ≥ 126

Comments

Primary sample: Plasma
Methodology :Enzymatic reference method with hexokinase

Thyroid Stimulating Hormone (TSH)	3.02	uIU/mL	0.27 - 4.2
-----------------------------------	------	--------	------------

Comments

Note: TSH Reference range in pregnant women is varied according to the gestational age:

- First trimester: 0.100-2.500 uIU/mL
- Second trimester: 0.200-3.000 uIU/mL
- Third trimester: 0.300-3.000 uIU/mL

Primary sample: Serum
Methodology: Electrochemiluminescence

Reviewed By:



Dr. Shailendra Rathod.MD
Laboratory Director
Physician Specialist Clinical
Pathology
License no. 10944644-001

** End Of Report **

Verified By : Shailendra Rathod.MD

Printed By: System

All Rights Reserved © National Technology

Printed Date PM 14:44 21/08/2022

Page 1 Of 1

Powered by LDM www.nt-me.com