



BML545792

Laboratory Investigation Report

Name	: Mr. JOSHUA LUKE SIMPSON	Ref No.	: 46379
DOB	: 22/06/2012	Sample No.	: 2504559308
Age / Gender	: 12 Y / Male	Collected	: 05/04/2025 20:00
Referred by	: Dr. AMAIZAH	Registered	: 06/04/2025 15:01
Centre	: CITICARE MEDICAL CENTER	Reported	: 06/04/2025 15:47

BIOCHEMISTRY

Test	Result	Flag	Unit	Reference Range	Methodology
C-REACTIVE PROTEIN (CRP)	1.4		mg/L	< 5.0 Please note change. Source: Roche IFU.	Particle-enhanced immunoturbidimetric assay

INTERPRETATION NOTES :

- CRP measurements are used as aid in diagnosis, monitoring, prognosis, and management of suspected inflammatory disorders and associated diseases, acute infections and tissue injury.
- C-reactive protein is the classic acute phase protein in inflammatory reactions.
- CRP is the most sensitive of the acute phase reactants and its concentration increases rapidly during inflammatory processes. The CRP response frequently precedes clinical symptoms, including fever. After onset of an acute phase response, the serum CRP concentration rises rapidly and extensively. The increase begins within 6 to 12 hours and the peak value is reached within 24 to 48 hours. Levels above 100 mg/L are associated with severe stimuli such as major trauma and severe infection (sepsis).
- CRP response may be less pronounced in patients suffering from liver disease.
- CRP assays are used to detect systemic inflammatory processes (apart from certain types of inflammation such as systemic lupus erythematosus (SLE) and Colitis ulcerosa); to assess treatment of bacterial infections with antibiotics; to detect intrauterine infections with concomitant premature amniorrhexis; to differentiate between active and inactive forms of disease with concurrent infection, e.g. in patients suffering from SLE or Colitis ulcerosa; to therapeutically monitor rheumatic disease and assess anti-inflammatory therapy; to determine the presence of post-operative complications at an early stage, such as infected wounds, thrombosis and pneumonia, and to distinguish between infection and bone marrow transplant rejection.

Sample Type : Serum

End of Report

Dr. Adley Mark Fernandes
M.D (Pathology)
Pathologist

Gyoma V. Shah
Dr. Vyoma V Shah
M.D (Pathology)
Clinical Pathologist

Greeshma P Sidharthan
Greeshma P Sidharthan

This is an electronically authenticated report

Page 1 of 1

Printed on: 06/04/2025 15:49

Test result pertains only to the sample tested and to be interpreted in the light of clinical history. These tests are accredited under ISO 15189:2012 unless specified by (^). Test marked with # is performed in an accredited referral laboratory.

