



Patient Name : Ms. PELANGINE YAWEH NDULA Sample UID No. : 4088583

 Age / Gender
 : 28 Y / Female
 Sample Collected On
 : 19-06-2025 14:45

 Patient ID
 : QLD088396
 Registered On
 : 19-06-2025 17:17

 Referred By
 : Dr. AMAIZAH
 Reported on
 : 20-06-2025 00:03

Referral Client : CITICARE MEDICAL CENTER(INSURANCE) External Patient ID : 42898
Emirates ID / Passport No : Print Version : V.1

Department of BIOCHEMISTRY

<u>Investigation</u> <u>Results</u> <u>Flag</u> <u>Units</u> <u>Biological Reference Interval</u> <u>Method</u>

* C-REACTIVE PROTEIN (CRP) 23.3 H mg/L <5 Particle enhanced

immunoturbidimetric assay

Sample: Serum Comments:

CLINICAL IMPLICATIONS:

- 1. CRP is the most sensitive acute phase reactant that can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgeryor neoplastic proliferation. CRP levels may predict future cardiovascular events and can be used as a screening tool.
- 2. The traditional test of CRP has added significance over the elevated ESR, which may be influenced by altered physiologic states. CRP tends to increase before rises in antibody titres and ESR level occurs. CRP levels also tend to decrease sooner than ESR levels.
- 3. The traditional test for CRP is elevated in rheumatic fever, RA, myocardial infarction, malignancy, bacterial and viral infections. The positive test indicates active inflammation but not its cause. In RA, the traditional test for CRP becomes negative with successful treatment and indicates that the inflammation has subsided.
- 4.High sensitive measurement of CRP (hs-CRP) are useful in assessing vascular inflammation and cardiovascular stratification. A single test for hs-CRP may not reflect an individual patient basal hs-CRP level, therefore follow up tests or serial measurements may be required in patients presenting with increased hs-CRP levels.

INTERFERING FACTORS: Haemolysed or lipemic sample may alter the results.

REFERENCE:

- 1) Manual of Laboratory and Diagnostics -Frances Fischbach Marshall B. Dunning III [9th Edition]
- 2) Tietz clinical guide to Laboratory tests(Fourth edition) ALAN H.B.WU

- END OF REPORT -

Note:

"The analytes with asterix (*) symbol are non-accredited parameters.".
"QLabs compliance with ISO 15189:2022 standards"

Ebin C Lorance Lab Technologist

DHA No. 57146854-002



Dr. Vidhya Mohan Specialist Clinical Pathologist Clinical Pathologist DHA No. 23553203-004

Dr. Dheepa Manoharan Medical Director Specialist Microbiologist DHA No. 00231751-004

Page 1 of 2





Patient Name Sample UID No. : Ms. PELANGINE YAWEH NDULA : EB4088583

Age / Gender : 28 Y / Female **Sample Collected On** : 19-06-2025 14:45 Patient ID : QLD088396 Registered On : 19-06-2025 17:17 : 20-06-2025 06:51 Referred By Reported on : Dr. AMAIZAH

: 42898 **Referral Client** : CITICARE MEDICAL CENTER(INSURANCE) **External Patient ID** Emirates ID / Passport No :

Print Version : V.1

Department of HEMATOLOGY

COMPREHENSIVE COMPLETE BLOOD COUNT

<u>Investigation</u>	<u>Results</u>	<u>Flag</u>	<u>Units</u>	Biological Reference Interval	<u>Method</u>
HEMOGLOBIN	9.9	L	g/dl	12-15	photometric
RBC COUNT	4.57		10^6/uL	3.8-4.8	Electrical Impedance
HEMATOCRIT	30.8	L	%	37-47	Calculation
MCV	67.3	L	fL	78-100	Calculation
МСН	21.6	L	pg	27-31	Calculation
МСНС	32.2		g/dl	31-35	Calculation
RDW	18	Н	%	9.3-16	Calculation
RDW-SD	42		fL	38.9-49	Calculation
MPV	8.4	L	fL	8.8-12.5	Calculation
PLATELET COUNT	318		10^3/uL	150-400	Electrical Impedance
* PCT	0.3		%	0.01-9.99	Calculation
* PDW	16.1			0.1-99.9	Calculation
* NUCLEATED RBC (NRBC)^	0.45		/100 WBC		Flow Cytometry
* ABSOLUTE NRBC COUNT^	0.03		10^3/uL		Calculation
* EARLY GRANULOCYTE COUNT (EGC)^	0.83		%		Flow Cytometry
* ABSOLUTE EGC^	0.05		10^3/uL		Calculation
WBC COUNT	6.5		10^3/uL	4-11	Electrical Impedance
* Neutrophil	75.25		%	40-80	VCS-Method
* Lymphocyte	7.56	L	%	20-40	VCS-Method
* Eosinophil	0.22	L	%	1-8	VCS-Method
* Monocyte	16.21	н	%	2-10	VCS-Method
* Basophil	0.76		%	0-2	VCS-Method
* ABSOLUTE NEUTROPHIL COUNT	4.87		10^3/uL	1.5-7	Calculation
* ABSOLUTE LYMPHOCYTE COUNT	0.49	L	10^3/uL	1.5-4	Calculation
* ABSOLUTE MONOCYTE COUNT	1.05	н	10^3/uL	0-0.8	Calculation
* ABSOLUTE EOSINOPHIL COUNT	0.01		10^3/uL	0-0.6	Calculation
* ABSOLUTE BASOPHIL COUNT	0.05		10^3/uL	0-0.2	Calculation
Sample: EDTA Whole Blood					

Sample: EDTA Whole Blood

- END OF REPORT -

Note:

"The analytes with asterix (*) symbol are non-accredited parameters.". "QLabs compliance with ISO 15189:2022 standards"

> **Ebin C Lorance** Lab Technologist



DHA No. 57146854-002

Dr. Vidhya Mohan **Specialist Clinical Pathologist Clinical Pathologist** DHA No. 23553203-004

Page 2 of 2