



Patient Name : Mr. EL MAHJOUB BOURHIM Sample UID No. : 4091839

 Age / Gender
 : 34 Y / Male
 Sample Collected On
 : 30-06-2025 16:50

 Patient ID
 : QLD091639
 Registered On
 : 30-06-2025 16:53

 Referred By
 : DR FARHAN
 Reported on
 : 30-06-2025 22:08

Referral Client : CITICARE MEDICAL CENTER(INSURANCE) External Patient ID : 47289
Emirates ID / Passport No : Print Version : V.1

Department of BIOCHEMISTRY

<u>Investigation</u> <u>Results</u> <u>Flag</u> <u>Units</u> <u>Biological Reference Interval</u> <u>Method</u>

* C-REACTIVE PROTEIN (CRP) 0.6 mg/L <5 Particle enhanced

immunoturbidimetric assay

Sample: Serum Comments:

CLINICAL IMPLICATIONS:

- 1. CRP is the most sensitive acute phase reactant that can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgeryor neoplastic proliferation. CRP levels may predict future cardiovascular events and can be used as a screening tool.
- 2. The traditional test of CRP has added significance over the elevated ESR, which may be influenced by altered physiologic states. CRP tends to increase before rises in antibody titres and ESR level occurs. CRP levels also tend to decrease sooner than ESR levels.
- 3. The traditional test for CRP is elevated in rheumatic fever, RA, myocardial infarction, malignancy, bacterial and viral infections. The positive test indicates active inflammation but not its cause. In RA, the traditional test for CRP becomes negative with successful treatment and indicates that the inflammation has subsided.
- 4.High sensitive measurement of CRP (hs-CRP) are useful in assessing vascular inflammation and cardiovascular stratification. A single test for hs-CRP may not reflect an individual patient basal hs-CRP level, therefore follow up tests or serial measurements may be required in patients presenting with increased hs-CRP levels.

INTERFERING FACTORS: Haemolysed or lipemic sample may alter the results.

REFERENCE:

- 1) Manual of Laboratory and Diagnostics -Frances Fischbach Marshall B. Dunning III [9th Edition]
- 2) Tietz clinical guide to Laboratory tests(Fourth edition) ALAN H.B.WU

- END OF REPORT -

Note:

"The analytes with asterix (*) symbol are non-accredited parameters.".
"QLabs compliance with ISO 15189:2022 standards"

Mojahid Babiker Lab Technologist

DHA No: 82488066-001



Dr. Vidhya Mohan Specialist Clinical Pathologist Clinical Pathologist DHA No. 23553203-004 Dr. Dheepa Manoharan Medical Director Specialist Microbiologist DHA No. 00231751-004

Page 1 of 2





Patient Name Sample UID No. : Mr. EL MAHJOUB BOURHIM : EB4091839

Age / Gender : 34 Y / Male $\textbf{Sample Collected On} \hspace{0.1in} \textbf{:} \hspace{0.1in} 30\text{-}06\text{-}2025 \hspace{0.1in} 16\text{:}50$ Patient ID : QLD091639 Registered On : 30-06-2025 16:53 : 30-06-2025 22:08 Referred By Reported on : DR FARHAN

: 47289 **Referral Client** CITICARE MEDICAL CENTER(INSURANCE) **External Patient ID** Emirates ID / Passport No : **Print Version** : V.1

Department of HEMATOLOGY

COMPREHENSIVE COMPLETE BLOOD COUNT

<u>Investigation</u>	<u>Results</u>	<u>Flag</u>	<u>Units</u>	Biological Reference Interval	<u>Method</u>
HEMOGLOBIN	14.6		g/dl	13-17	photometric
RBC COUNT	4.82		10^6/uL	4.5-5.5	Electrical Impedance
HEMATOCRIT	43.2		%	42-52	Calculation
MCV	89.6		fL	78-100	Calculation
МСН	30.4		pg	27-31	Calculation
мснс	33.9		g/dl	31-35	Calculation
RDW	13.3		%	9.3-16	Calculation
RDW-SD	42.9		fL	38.9-49	Calculation
MPV	11.2		fL	8.8-12.5	Calculation
PLATELET COUNT	188		10^3/uL	150-400	Electrical Impedance
* PCT	0.2		%	0.01-9.99	Calculation
* PDW	17.6			0.1-99.9	Calculation
* NUCLEATED RBC (NRBC)^	0.87		/100 WBC		Flow Cytometry
* ABSOLUTE NRBC COUNT^	0.07		10^3/uL		Calculation
* EARLY GRANULOCYTE COUNT (EGC)^	0.17		%		Flow Cytometry
* ABSOLUTE EGC^	0.01		10^3/uL		Calculation
WBC COUNT	7.7		10^3/uL	4-11	Electrical Impedance
* Neutrophil	75.1		%	40-80	VCS-Method
* Lymphocyte	16.94	L	%	20-40	VCS-Method
* Eosinophil	1.97		%	1-8	VCS-Method
* Monocyte	5.42		%	2-10	VCS-Method
* Basophil	0.57		%	0-2	VCS-Method
* ABSOLUTE NEUTROPHIL COUNT	5.76		10^3/uL	1.5-7	Calculation
* ABSOLUTE LYMPHOCYTE COUNT	1.3	L	10^3/uL	1.5-4	Calculation
* ABSOLUTE MONOCYTE COUNT	0.42		10^3/uL	0-0.8	Calculation
* ABSOLUTE EOSINOPHIL COUNT	0.15		10^3/uL	0-0.6	Calculation
* ABSOLUTE BASOPHIL COUNT	0.04		10^3/uL	0-0.2	Calculation
Sample: EDTA Whole Blood					

Sample: EDTA Whole Blood

- END OF REPORT -

Note:

"The analytes with asterix (*) symbol are non-accredited parameters.". "QLabs compliance with ISO 15189:2022 standards"

> Ebin C Lorance Lab Technologist



Dr. Vidhya Mohan **Specialist Clinical Pathologist Clinical Pathologist** DHA No. 23553203-004

Page 2 of 2